Diabetes During Pregnancy — Continued

TABLE 2. Number and rate* of diabetes for women born in the 50 states and the District of Columbia (DC) and for women born elsewhere, by race/ethnicity — United States, 1993-1995

| | Women born in 50 states and DC | | | Women born elsewhere | | |
|---------------------------|--------------------------------|-----------------|---------------|----------------------|-----------------|---------------|
| Race/Ethnicity | No.† | Unadjusted rate | Adjusted rate | No.† | Unadjusted rate | Adjusted rate |
| Non-Hispanic | | | | | | |
| White | 6,653,662 | 25.2 | 24.3 | 332,677 | 27.2 | 23.0 |
| Black | 1,618,276 | 21.2 | 26.6 | 143,659 | 39.5 | 33.4 |
| Hispanic | | | | | | |
| Mexican | 494,906 | 23.2 | 31.1 | 834,834 | 22.5 | 25.7 |
| Puerto Rican | 96,380 | 28.0 | 36.2 | 64,137 | 37.0 | 41.4 |
| Cuban | 11,945 | 23.0 | 24.3 | 23,181 | 25.8 | 21.4 |
| Central or South | | | | | | |
| American | 18,347 | 17.6 | 21.3 | 252,773 | 26.0 | 24.3 |
| American Indian/ | | | | | | |
| Alaskan Native | 104,322 | 44.0 | 53.0 | 4,442 | 43.0 | 42.1 |
| Asian/Pacific Islander | | | | | | |
| Chinese | 6.914 | 39.1 | 28.6 | 70,171 | 39.0 | 27.1 |
| Japanese | 12,175 | 35.3 | 27.7 | 13,681 | 19.3 | 15.6 |
| Hawaiian | 16,568 | 28.8 | 32.7 | 410 | § | 33.2 |
| Filipino | 13,771 | 26.8 | 29.9 | 74,566 | 42.2 | 32.0 |
| Asian Indian¶ | 3,627 | 38.3 | 34.0 | 27,841 | 58.5 | 50.3 |
| Korean [¶] | 844 | § | § | 24,023 | 19.1 | 16.1 |
| Samoan¶ | 1,845 | 15.2 | 17.7 | 3,005 | 32.3 | 31.0 |
| Vietnamese [¶] | 351 | § | § | 33,745 | 24.3 | 19.4 |
| Total** | 9,280,027 | 24.8 | 25.0 | 2,078,873 | 28.0 | 26.4 |

^{*}Per 1000 singleton live-born infants in specified population.

†Women for whom place of birth and diabetes status were reported.

§Numbers were too small for meaningful analysis.

†Data were available for seven states (California, Hawaii, Illinois, New Jersey, New York, Texas, and Washington).

^{**} Includes races other than those listed.